

Simultaneous multi-cancer detection and tissue of origin (TOO) localization using targeted bisulfite sequencing of plasma cell-free DNA (cfDNA)

Minetta C. Liu, MD¹

Geoffrey R. Oxnard, MD²; Eric A. Klein, MD³; Michael V. Seiden, MD⁴; Earl Hubbell, PhD⁵; Oliver Venn, DPhil⁵; Arash Jamshidi, PhD⁵; Nan Zhang, PhD⁵; John F. Beausang, PhD⁵; Samuel Gross, PhD⁵; Kathryn N. Kurtzman, MD⁵; Eric T. Fung, MD, PhD⁵; Jessica Yecies, PhD⁵; Rita Shaknovich, MD, PhD⁵; Alexander P. Fields, PhD⁵; Mikkael A. Sekeres, MD³; Donald Richards, MD, PhD⁶; Peter P. Yu, MD⁷; Alexander M. Aravanis, MD, PhD⁵; Anne-Renee Hartman, MD⁵

¹Mayo Clinic, Rochester, MN. ²Dana Farber Cancer Institute, Boston, MA. ³Cleveland Clinic, Cleveland, OH. ⁴US Oncology Research, The Woodlands, TX. ⁵GRAIL, Inc., Menlo Park, CA. ⁶Texas Oncology, Tyler, TX. ⁷Hartford HealthCare Cancer Institute, Hartford, CT.

Disclosures

Study funded by GRAIL, Inc. The Mayo Clinic was compensated for Minetta C. Liu's advisory board activities for GRAIL, Inc. GRO is an advisory board member and consultant for Inivata Ltd.; an honorarium recipient from Guardant Health, Inc., Sysmex Corporation, and Bio-Rad Laboratories, Inc.; and a consultant for DropWorks, Inc., AstraZeneca plc, and GRAIL, Inc. EAK is a consultant for GRAIL, Inc. and Cellanyx, LLC. MVS is an employee of, and shareholder in, McKesson Corporation. EH, OV, AJ, NZ, JFB, SG, KNK, ETF, JY, RS, APF, ANA, and A-RH are employees of GRAIL, Inc., with equity in the company. AJ and KNK hold stock in Illumina, Inc.


The Circulating Cell-free Genome Atlas (CCGA) Study: Supporting Development of a Multi-Cancer Test

Prospective, observational, longitudinal, case-control study for the discovery, training, and validation of a multi-cancer test



Targeted Methylation Assay for Further Development

Pre-specified studies for discovery and validation




Discovery phase completed in 2018^{1,2}

Targeted methylation was selected for further assay development, including training and internal cross-validation³

Here, we report on an independent validation set that is ongoing

Target Selection Using Machine Learning Algorithm

Targeted methylation panel developed through generation and analysis of an extensive database of plasma and tissue methylation patterns



*Upper GI combines esophageal and gastric cancers: diagnostic workup covers both cancer types.



Stage-specific Performance was Consistent between Training and Validation Sets


- Cross-validated training set was independently tested in the validation set
- Trained classifier targeted >99% specificity and achieved a consistent <1% FPR in the training and validation sets ($P=0.095$)
- Training and validation set detection rates were consistent across stages for the pre-specified cancers and all cancers
- Confirms training data were not overfitted



More than 20 Cancer types Detected at Early and Late Stages

At 99.3% specificity:


- Sensitivity overall was 55% (51-59%)
- Sensitivity in pre-specified[†] cancer types was 76% (72-81%)



*Plot excludes unstaged cancers. †Includes anal, bladder, colorectal, esophageal, head and neck, liver/bile-duct, lung, lymphoma, ovary, pancreatic, plasma cell neoplasm, stomach cancer..

Highly Accurate Tissue of Origin (TOO) Localization

- 96% of samples with assigned TOO
- 93% of those calls were correct
- Highly precise localization to a single tissue site across >20 distinct tumor types




*Upper GI combines esophageal and gastric cancers: diagnostic workup covers both cancer types.

**Other cancer types= skin cancer (not including basal cell carcinoma, squamous cell carcinoma, or melanoma), testis, seminoma, vagina, and vulva.

☰☰☰ Tissue of Origin (TOO) Accuracy is Consistently High Across Stages

- Single tissue localization: **93% of TOO calls were correct**
- Localization to the top two TOO calls: **95% of TOO calls were correct**



*Lymphoid leukemias, myeloid neoplasms, and brain. †Includes anal, bladder, colorectal, esophageal, head and neck, liver/bile-duct, lung, lymphoma, ovary, pancreatic, plasma cell neoplasm, stomach.

≡≡≡ Conclusions

- Multiple deadly cancer types that currently have no screening paradigm were detected across stages and simultaneously accurately localized to a TOO using methylation signatures in plasma cfDNA.
- This was achieved with trained thresholds that resulted in a single, fixed, low false positive rate ($<1\%$) in an independent validation set.
- Importantly, results in the independent validation set were indistinguishable from the training set, demonstrating the robustness of machine learning classifier training, with no evidence of overtraining.
- This validation demonstrates feasibility of a single blood-based test that can simultaneously detect multiple cancers and supports further clinical development to prepare for the return of results.



Acknowledgements

- Study participants who graciously donated their time, energy, and specimens
- CCGA investigators and collaborators for advice, participant enrollment, and collection of data and specimens
- Advisors and Scientific Advisory Board members for their helpful feedback and advice
- Tony Wu, Lori Zhang, and Hai Liu, PhD (GRAIL, Inc.) for statistical support and figure generation
- The many GRAIL teams who have worked and continue to work on this study